

2010. *Herpetol. Rev.* 41:376). This is Entomology Contribution No. 1186, DPI, FDACS.

**LINDA S. STEVENSON** (e-mail: stevenl@doacs.state.fl.us) and **LOUIS A. SOMMA**, Florida State Collection of Arthropods, Division of Plant Industry—Entomology Sect., Florida Department of Agriculture & Consumer Services, PO Box 147100, 1911 SW 34th Street, Gainesville, Florida 32614-7100, USA (e-mail: sommal@doacs.state.fl.us).

**PELOBATES VARALDII (Moroccan Spadefoot Toad)**. MOROCCO: temporary ponds next to road N1 between Asilah and Tanger (35.621983°N, 5.979220°W; WGS84). 4 April 2009. Marta Lapeña and Luis Javier Barbadillo. Verified by mt-DNA barcoding (IMS, unpubl. data). Tissue vouchers of five tadpoles deposited at Instituto de Investigación en Recursos Cinegéticos (IMS personal collection, voucher numbers: IMS 1148–1152). Hundreds of tadpoles in three ponds < 100 m of each other, in all cases co-occurring with larvae of *Pleurodeles waltl*. Extends range 50 km N (town of Larache; see Bons and Geniez 1996. *Anfibios y Reptiles de Marruecos* [incluido Sáhara Occidental]. Atlas Biogeográfico. Asociación Herpetológica Española, Barcelona. 320 pp.).

**MARTA LAPEÑA**, Avenida del Ejército Español, 1, Bloque 1, 6° D, 51002 Ceuta, Spain (e-mail: martalape@yahoo.es); **LUIS JAVIER BARBADILLO**, Avenida Hassan II, 2, 93000 Tetouan, Morocco (e-mail: ljbarbadillo@yahoo.es); **ÍÑIGO MARTÍNEZ-SOLANO**, Instituto de Investigación en Recursos Cinegéticos, Ronda de Toledo, s/n, 13005 Ciudad Real, Spain (e-mail: inigomsolano@irec.csic.es).

**PROCERATOPHRYS PAVIOTII**. BRAZIL: ESPÍRITO SANTO: Pinheiros, Fazenda Santa Rita (18.30°S, 40.15°W; WGS 84; 105 m elev.). 25 August 2008. J. Peres and J. E. Simon. Museu de Biologia Professor Mello Leitão, Santa Teresa, Espírito Santo, Brazil (MBML 7237–7243). Verified by G. Machado Prado. The species was previously known only from its type locality: Estação Biológica de Santa Lúcia (ca. 19.96°S, 40.53°W, elev. 560 m), Municipality of Santa Teresa, mountainous region of the State of Espírito Santo, southeastern Brazil (Cruz et al. 2005. *Arq. Mus. Nac.* 63[2]:289–295). This record extends the species distribution ca. 190 km N from type locality.

**JULIANA PERES** and **JOSÉ EDUARDO SIMON**, Museu de Biologia Prof. Mello Leitão, Rua José Ruschi, 4, Santa Teresa, ES, Brazil, CEP: 29650-000 (e-mail: simon@ebr.com.br)

**PSEUDACRIS CRUCIFER (Spring Peeper)**. USA: GEORGIA: MERTHWETHER Co.: Joe Kurz Wildlife Management Area, 0.64 km S of Damascus Church Rd (33.12956°N, 84.53502°W, WGS 84; elev. 237 m). 11 November 2009. Robert L. Hill. Verified by Craig Guyer. AUM AHAP-D 284. New county record (Jensen et al. 2008. *Amphibians and Reptiles of Georgia*. Univ. of Georgia Press, Athens. 575 pp.); previously documented in all surrounding counties. Specimen found under decaying log, photographed, and released.

**ROBERT L. HILL**, Department of Conservation and Research, Atlanta Botanical Garden, Atlanta, Georgia 30309, USA; e-mail: rhill@atlantabotanicalgarden.org.

**PSEUDACRIS CRUCIFER (Spring Peeper)**. USA: TENNESSEE: UNION Co.: Maynardville (36.219444°N, 83.828889°W; WGS84). 27 September 2010. Stephen K. Nelson. Verified by A. Floyd Scott. David H. Snyder Museum of Zoology, Austin Peay State University (APSU 19063, color photo). First county record (Redmond and Scott 1996. *Atlas of Amphibians in Tennessee*. Misc. Publ. No. 12, Center for Field Biology, Austin Peay State University, 24 pp. Internet version, available at <http://www1.apsu.edu/amatlas/index.html>. Accessed 25 October 2010). On road at night in rain, at an air temperature of 15°C.

**STEPHEN K. NELSON**, University of Tennessee, Knoxville, Tennessee 37996-0925, USA; e-mail: snelson24@utk.edu

**PSEUDACRIS FERIARUM (Upland Chorus Frog)**. USA: TENNESSEE: MARSHALL Co.: (35.6700°N, 86.7856°W; NAD83). 21 April 2009. Michael Anderson and Daniel Estabrooks. Verified by A. Floyd Scott. Austin Peay State University's David H. Snyder Museum of Zoology (APSU 19059, photographic voucher). First record for county, ca. 38 km from previous site in Giles Co. (Redmond and Scott 1996. *Atlas of Amphibians in Tennessee*. Misc. Publ. No. 12. The Center for Field Biology, Austin Peay State University, Clarksville, Tennessee. 94 pp. Internet version [<http://www.apsu.edu/amatlas>] contains links to information regarding Tennessee distribution of amphibians recorded since 1996; accessed 07 October 2010). Male and female in amplexus.

**MICHAEL A. ANDERSON** (e-mail: maa3t@mtmail.mtsu.edu) and **DANIEL B. ESTABROOKS** (e-mail: dbe2g@mtmail.mtsu.edu), Department of Biology, Middle Tennessee State University, Murfreesboro, Tennessee 37132, USA.

**RHINELLA BERNARDOI**. ARGENTINA: SAN JUAN: DEPARTAMENTO CAUCETE: Quebrada de Las Flores (31.70083°S, 68.09500°W, WGS 84; elev. 743.5 m). February 2009. L. Quiroga, E. Sanabria, F. Cano, R. Cortez, and V. Marinero. Herpetological Collection, Fundación Miguel Lillo, Tucumán, Argentina (FML 23921, male, SVL 70 mm, body mass 40 g, and body temperature 18°C). Verified by F. Arias.

Previously known only from the type locality, this record extends known distribution about 170 km S from Parque Provincial Ischigualasto, Departamento Valle Fértil, San Juan, Argentina (Sanabria et al. 2010. *Zootaxa* 2396:50–60).

**EDUARDO A. SANABRIA** (e-mail: sanabria.eduardoa@gmail.com), **LORENA B. QUIROGA**, **FLORENCIA A. CANO**, **RICARDO O. CORTEZ**, and **NANCY V. MARINERO**, Departamento de Biología, F.C.E.F. y N., Universidad Nacional de San Juan, Avenida España 400 (N) C.P. 5400, San Juan, Argentina.

**RHINELLA RUBESCENS (Rufous Toad)**. BRAZIL: MATO GROSSO DO SUL: MUNICIPALITY OF DOURADOS: (22.09613°S, 55.37456°W; SAD 69). 16 June 2007. C. Aoki and P. Landgraf Filho. Coleção Zoológica de Referência da Universidade Federal de Mato Grosso do Sul, Mato Grosso do Sul, Brazil (ZUFMS-AMP 1213). Verified by R. A. Brandão. First record for Mato Grosso do Sul State (Frost 2010. *Amphibian Species of the World: an Online Reference*. Version 5.4). The geographic distribution of *Rhinella rubescens* encompasses the Brazilian states of Pará, Piauí, Goiás, Minas Gerais, São Paulo, and Bahia. Extends the known distribution of *R. rubescens* ca. 520 km SW from the closest known locality (Goiás).

**CAMILA AOKI**, Programa de Pós Graduação em Ecologia e Conservação. Universidade Federal de Mato Grosso do Sul, Cidade Universitária, s/n, Bairro Universitário, CEP 79070-900, Campo Grande, MS, Brazil (e-mail: aokicamila@yahoo.com.br); **PAULO LANDGRAF FILHO**, Centro de Ciências Biológicas e da Saúde, Universidade Federal de Mato Grosso do Sul, Laboratório de Zoologia, CEP 79070-900, Campo Grande, MS, Brazil; **FABRÍCIO HIROIUKI ODA**, Programade Pós-Graduação em Ecologia de Ambientes Aquáticos Continentais. Universidade Estadual de Maringá, Nupélia - Núcleo de Pesquisas em Limnologia, Ictiologia e Aqüicultura, Laboratório de Ictioparasitologia – Bloco G-90, Av. Colombo, 5790, CEP 87020-900, Maringá, PR, Brazil; **ROBERTO MACEDO GAMARRA**, Programa de Pós Graduação em Ecologia e Conservação. Universidade Federal de Mato Grosso do Sul, Cidade Universitária, s/n, Bairro Universitário, CEP 79070-900. Campo Grande, MS, Brazil.

**SCINAX AURATUS (Golden Tree Frog)**. BRAZIL: PARAÍBA: Manguape, Reserva Biológica Guaribas (06.74339°S, 035.14711°W, WGS84; elev. 182 m). 5 June 2010. Collected by A. Santos Protázio,

L. B. Q. Cavalcanti. Verified by G. H. Calazans Vieira. Departamento de Sistemática e Ecologia, Coleção Herpetológica, Universidade Federal da Paraíba (SVL 22.38 mm; UFPB 4884; collecting license ICMBIO 15806-2). First state record. *Scinax auratus* was previously known from the states of Bahia, Sergipe, and Alagoas (Alves et al. 2004. Rev. Brasil. Zool. 21[2]:315–317; Bokermann 1969. Rev. Brasil. Biol. 29[2]:159–162; Juncá 2005. In Juncá et al. [org.], Biodiversidade e Conservação da Chapada Diamantina, pp. 337–356. Ministério do Meio Ambiente; Nunes and Pombal Jr. 2010. Amphibia-Reptilia 31:347–353; Santana et al. 2009. Check List 5[2]:246–248). Extends the species distribution ca. 406 km NE from the municipality of Quebrângulo, Alagoas state.

**ARIELSON DOS SANTOS PROTÁZIO, LUCAS BARBOSA DE QUEIROGA CAVALCANTI, DANIEL OLIVEIRA MESQUITA**, Universidade Federal da Paraíba, Departamento de Sistemática e Ecologia, Cidade Universitária, Cep: 58059-900, João Pessoa, Paraíba, Brazil; **AIRAN DOS SANTOS PROTÁZIO**, Universidade Estadual de Feira de Santana, Laboratório de Animais Peçonhentos e Herpetologia, Av. Transnordestina, S/N, Novo Horizonte, Cep: 44036-900, Feira de Santana, Bahia, Brazil (e-mail: neu\_ptz@hotmail.com).

**SCINAX GRANULATUS (NCN)**. ARGENTINA: BUENOS AIRES: PARTIDO DE GENERAL PUEYRREDON: Sierra de Los Padres (37.94028°S, 57.79472°W; WGS 84). 24 July 2010. L. E. Vega and P. J. Bellagamba. Verified by J. Faivovich. Herpetological Collection of Laboratorio Vertebrados, Departamento de Biología, Universidad Nacional de Mar del Plata, Mar del Plata, Buenos Aires, Argentina (UNMDP 1809–1810). First citation for the county and the southernmost record known for this species. This is the first documentation of this frog inhabiting mountain environments at 170 m elev. Present record extends the distribution of this frog 120 km to the SW from nearest known locality in Partido de General Madariaga in the Buenos Aires province (Barrasso and Cajade 2007. Herpetol. Rev. 38:99). The species was previously known from northeastern Argentina, south to Buenos Aires Province; Uruguay and Paraná, Santa Catarina, and Rio Grande do Sul, Brazil; expected in southern Paraguay (Frost 2010. Amphibian Species of the World: an Online Reference. Version 5.4 [8 April 2010]). Electronic database accessible at <http://research.amnh.org/vz/herpetology/amphibia>. American Museum of Natural History, New York). Barrasso and Cajade (2007, *op. cit.*) pointed out that Gallardo mentioned two specimens from Sierra de la Ventana, but the citation is not supported by voucher specimens. Two adults were captured under rocks. Another five individuals were found inactive under rocks.

**LAURA E. VEGA, PATRICIO J. BELLAGAMBA, OSCAR A. STELLATELLI**, Laboratorio Vertebrados, Departamento de Biología, Universidad Nacional de Mar del Plata, Funes 3250, 7600 Mar del Plata, Argentina; **GUILLERMO S. NATALE**, Centro de Investigaciones del Medio Ambiente, Departamento de Química, Facultad de Ciencias Exactas, Universidad de La Plata, Calle 47 y 115 s/n, La Plata, Buenos Aires, Argentina.

**SCINAX TRAPICHEIROI**. BRAZIL: RIO DE JANEIRO: MUNICIPALITY OF BOM JESUS DO ITABAPOANA: River Itabapoana bank (21.106424°S, 41.725598°W; SAD 69, elev. 139 m). 18 August 2006. Collected by A. L. Silveira. Museu Nacional / UFRJ, Rio de Janeiro, RJ, Brazil (MNRJ 42510). Verified by U. Caramaschi. The new record extends the known geographic distribution of species about 205 km NE from the closest previous record (Saquarema, ZUFJR 6720–6723, Luna-Dias et al. 2009. Check List 5[2]:251–253), and constitutes the northern and western boundaries of this distribution. First record of *Scinax trapicheiroi* in the Atlantic Forest biome, a semideciduous seasonal forest. Species was previously known from municipalities of Rio de Janeiro, Maricá, Angra dos Reis (Ilha Grande),

Mangaratiba (Ilha da Marambaia), Pirai and Saquarema, in Rio de Janeiro state, Brazil (Luna-Dias et al., *op. cit.*; Lutz 1954. Mem. Inst. Oswaldo Cruz 52[1]:155–197); all localities occur in area of dense ombrophile forest.

**ADRIANO LIMA SILVEIRA**, Setor de Herpetologia, Departamento de Vertebrados, Museu Nacional / Universidade Federal do Rio de Janeiro, Quinta da Boa Vista, São Cristóvão, CEP 20940-040, Rio de Janeiro, RJ, Brazil; e-mail: biosilveira@yahoo.com.br.

## CROCODYLIA — CROCODILIANS

**PALEOSUCHUS PALPEBROSUS (Cuvier's Dwarf Caiman, Jacaré-paguá)**. BRAZIL: CEARÁ: ITAPIPOCA MUNICIPALITY: Sítio São José (03.1825°S, 39.4369444°W; WGS84). January–February 2005. F. H. C. de Lima. Collection of Herpetology of Núcleo Regional de Ofiologia (NUROF) - Universidade Federal do Ceará, Fortaleza, Brazil (CHUFC C045–C046). Verified by L. M. Verdade. First state record. The species was also photographed and registered by visual survey on Serra das Almas, Crateús Municipality (03.345°S, 40.8861111°W; datum WGS84), located 260 km airline SW from Itapipoca, during January 2003. The species was previously known from basins of Amazonian (Colombia, Venezuela, Guianas, Bolívia, Paraguay and Brazil), Pantanal, Caatinga, and Atlantic Forest (Brazil) biomes (Carvalho 1951. Os Jacarés do Brasil. Arquivos do Museu Nacional, XLII:127–152; King and Burke 1989. Crocodilian, Tuatara, and Turtle Species of the World. Association of Systematics Collections, Washington, DC. 216 pp). In northeastern Brazil, the species was known from Grande River in the São Francisco River basin, state of Bahia (Carvalho, *op. cit.*), and recently from Atlantic Forest on João Pessoa Municipality, state of Paraíba (Santana et al. 2008. Biotemas 21[1]:75–84). Extends known distribution ca. 660 km airline NW from João Pessoa, and ca. 900 km airline SW of previously known range to the Rio Grande/Rio São Francisco. This is the first record of the species from streams in dune forests (Complexo Litorâneo), ca. 3 km from the seacoast, and is the northernmost distribution point in northeastern Brazil.

**DANIEL CASSIANO LIMA**, Programa de Pós Graduação em Biodiversidade Animal (PPGBA/UFESM), Cidade Universitária, Camobi, Km 9, 97105-700, Santa Maria, Rio Grande do Sul, Brazil; Universidade Estadual do Ceará, Av Mons. Tabosa s/n, 62500-000, Itapipoca, Ceará, Brazil; Doctorate fellowship of Fundação Cearense de Apoio ao Desenvolvimento Científico e Tecnológico-FUNCAP (e-mail: dancassiano@yahoo.com.br); **FRANCISCO HÉLIO COELHO DE LIMA**, Programa de Pós-Graduação em Desenvolvimento e Meio Ambiente (PRODEMA/UFC), Campus do Pici, Bloco 902, 60455-760, Fortaleza, Ceará, Brazil; Master Fellowship of Coordenação de Aperfeiçoamento de pessoal de Nível Superior-CAPES (e-mail: heliosbio74@yahoo.com.br); **DIVA MARIA BORGES-NOJOSA**, Universidade Federal do Ceará, NUROF-UFC, Campus do Pici, Bloco 905, 60455-760, Fortaleza, Ceará, Brazil (e-mail: dmbnojosa@yahoo.com.br).

## TESTUDINES — TURTLES

**APALONE SPINIFERA (Spiny Softshell)**. USA: OHIO: UNION Co.: Paris Township: Mill Creek at Swartzkopf Park in Marysville (40.24449°N, 83.37324°W; WGS 84). 16 July 2008. Jeffrey E. Dawson. Verified by Jeffrey G. Davis. Photo voucher in Cincinnati Museum Center, Geier Collections and Research Center (CMC HP 480). First vouchered record for the county (Wynn and Moody 2006. Ohio Turtle, Lizard, and Snake Atlas. Ohio Biol. Surv. Misc. Contr. No. 10, Columbus. iv + 81 pp.). Several specimens photographed as they basked on large woody debris within the stream. Rausch (1947. Am. Midl. Nat. 38:434–442) reported obtaining one *A. spinifera*